



QRG – Thromboprophylaxis for inpatients (adults)				
No.	Action	Responsibility		
1.	There are two ways of ordering mechanical or pharmaceutical Thromboprophylaxis for inpatients.	Doctor / NMP		
	The preferred way (and easier as VTE assessment and VTE prophylaxis prescribing can be done from one screen) is the from the VTE Risk Assessment form using the VTE Prophylaxis Care Set . – Continue with Steps 2-7 .			
	Alternatively Thromboprophylaxis can be ordered from Requests / Care plans using the Heparin / LMWH Powerplan (continue with step 8). This may be suitable if VTE Risk Assessment has already been completed but a change in clinical status may require a change in prescribing.			
2.	In the VTE Risk Assessment form, from the Prescribing Tab click on +Add and search for "VTE prophylaxis"	Doctor / NMP		
	VTE Risk Assessment - ZZZEPMA, PHARM01			
	*Performed on: 05/01/2018 • 1422 • GMT			
	ZZZEPMA, PHARM01 Prescribing	9		
	Postnatal Recommendations			
	* surg/Med Based on your VTE Risk Assessment Prescribe Prophylaxis where appropriate Prescribing			
	as the order set.	St		
	+ Add Document Medication by History Reconciliation - Sheck Interactions (Or Medication Sheck Sheck Interactions) (Or Medication Sheck Sheck Sheck Interactions)	Sta Y		
	View Display: All Active Medications, All Inactive Medications 24 Hrs Back 🗸			
	Orders for Signature	Det		
	On Categorised Diet Ordered	00		
	Click on the VTE Prophylaxis Care set to open the set.			
	Search: vte prophylaxis 🔍 🍕 💘 Advanced Options 👻 Type: 👘 Inpatient			
	Tenter to Search Sepsis Search within:			



Milton Keynes University Hospital NHS Foundation Trust

3.	Select the required orders by ticking the box next to it, e.g. Antiembolic stockings and Dalteparin 5000 Units for a patient weighing 46-100kg. If Heparin is indicated for prophylaxis (NOT treatment) then this can be ticked from this care set as well.	Doctor / NMP
4.	Du Table OK Cancel	Doctor / NMP
	Click	
5.	Click Done	Doctor / NMP
6.	Click	Doctor / NMP
7.	Ensure all other mandatory fields have been completed in the form and then click to sign the form. Then proceed to step 15.	Doctor / NMP
8.	If Thromboprophylaxis is ordered outside a VTE Risk Assessment form, this can be done from the Request / Care Plans tab in PowerChart. Click on +Add and type in "heparin"	Doctor / NMP
	Search: heparin 🔍 📆 Advanced Options 👻 Type: 👘 In	
	📭 🖆 🚖 🔹 🖆 Folder: Search within: 📶	
	Heparin Heparin / Low Molecular Weight Heparin (LMWH) Care Plan	
	IMPORTANT : If different Thromboprophylaxis is ordered via this step, please note that the original order still needs to be cancelled to avoid duplication.	
9.	Click on the Powerplan and select Done to open the plan	Doctor / NMP
10.	In the plan navigate to the section for prophylaxis of thromboembolisms Heparin / Low Molecular Weight Heparin (LMWH) Care Plan (Initiated Pending) Petert Care Point	Doctor / NMP





11.	Select the required orders by ticking the box next to them.	Doctor / NMP
	For Dalteparin tick the box and then select an appropriate order sentence. If a patient's weight has been documented, the weight filter for order sentences will only display the appropriate dose of Dalteparin based on the documented weight. If no weight has been documented, then all available order sentences will show.	
	Ø ∇ Component Status Details Heaparin (LNW/H) Care Plan (Initiated Pending)	
	A Patient Care See intranet for most up-to-date policies "Anticoaquilation Prescribing Guidance", "Thromboprophylaxis in Pregnancy and Puerperium"	
	COUR times a day FOUR times a day. Check that stockings are being worn according to Trust policy. Stockings must be removed on a daily basis so that the patient's legs can be was the skin can be assessed	hed
	d Medications For Prophylaxis of Thromboembolic event	
	K in Dateparin Select an order sentence Accumulation of LMWH can occur in renal impairment. For DOSE 2,500 unit - ROUTE: subcUTANEOUS - injection - ONCE a day (evening) - Thromboprophylaxis Greater Than or Equal To 18 year and Less	har
	Generation of the second	00 k 140
	Contractioner of measurement of measurement of measurement of the same time as the DoSE 5,000 unit - ROUTE subCUTANEOUS - injection - TWICE a day - for: Thromboprophylaxis and the same time as the DOSE 5,000 unit - ROUTE subCUTANEOUS - injection - ONCE a day (evening) - Prophylaxis in pregnancy, crCl < 30mL/min Less Than 50 kg	180
	DOSE 5,000 unit - ROUTE: subcUTTANEOUS - injection - TWICE a day - Thromboprophylaxis in pregnancy 50 - 90 kg DOSE 5,000 unit - ROUTE: subcUTANEOUS - injection - ONCE a day (evening) - Thromboprophylaxis in pregnancy 50 - 90 kg	
	Construction of the second secon	
	DOSE: 75 unit/kg - ROUTE: sub-CUTANEOUS - injection - ONCE a day (evening) - Thromboprophylaxis in pregnancy Greater Than or Equal To 170 kg	_
12.	Alternatively, if unfractionated Heparin is to be ordered, please tick the corresponding order from the prophylaxis section.	Doctor / NMP
13		Doctor / NMP
15.	Orders For Signature	
	Click	
14		Doctor / NMP
17.	Sign	
15	Navigate to drug chart to ensure all medication orders display correctly	Doctor / NMP
10.	having a to a ray shart to choose an incultation oracio display correctly.	